

WHICH RADIUS FOR THE SUN?

V. Castellani,¹ S. Degl'Innocenti,^{1,2} G. Fiorentini^{2,3}

¹ Dipartimento di Fisica, Università di Pisa, piazza Torricelli 2, I-56126 Pisa, Italy

² Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara, via Paradiso 12, I-44100 Ferrara, Italy

³ Dipartimento di Fisica, Università di Ferrara, via Paradiso 12, I-44100 Ferrara, Italy

Abstract

The high accuracy reached by solar limb observations, by helioseismic measurements and by Standard Solar Models (SSMs) calculations suggests that general relativity corrections are included when discussing the solar radius. The Allen value ($R_{\odot} = 695.99 \pm 0.07$ Mm) has to be reduced by 1.5 Km. This correction, which is small as compared with present accuracy, should be kept in mind for future more precise measurements and/or calculations.

Following a well established procedure, the observation of the solar disc combined with precise estimates of the astronomical unit has early produced high quality evaluations of the solar radius. According to Allen (1976) one finds:

$$R_{\odot} = 695.99 \pm 0.07 \text{ Mm}; \quad (\Delta R/R)_{\odot} = \pm 1 \cdot 10^{-4}. \quad (1)$$

As recently reviewed by Schou et al. (1997) and Brown and Christensen-Dalsgaard (1998), more recent estimates appear in substantial agreement with such a value. Thus, for a long time this value has been used to constrain Standard Solar Models with a precision largely exceeding the common requirements for stellar evolutionary models.

However, in more recent time helioseismology (HeS) has proved to be able to test the solar structure to an extraordinary degree of precision, comparable with the error quoted above, see e.g. Basu and Christensen-Dalsgaard (1997), Bahcall, Basu and Kumar (1997), Degl'Innocenti et al. (1997), Dziembowski (1996), Dziembowski et al. (1994). As a consequence, recent works (Schou et al., 1997, Basu, 1998, and Antia, 1998) discussed the possibility that helioseismology indicates a slightly smaller radius for the Sun, decreasing the above quoted value by a quantity

$$\delta R_{HeS} \approx 0.3 \text{ Mm}; \quad (\delta R/R)_{HeS} \approx 4 \cdot 10^{-4} \quad (2)$$

which, though rather small, is definitely beyond the formal error for Allen's value.

The issue has been recently discussed by Brown and Christensen-Dalsgaard (1998). They reanalyzed the already known inconsistency between the observational radius, $R_{\odot}$, defined as the radial distance to the inflection point of the limb intensity profile, and the photospheric radius R_{ph} , adopted by theoretical models, which is the point at the basis of the atmosphere with a temperature T equal to the star effective temperature, i.e., the point such that $R_{ph}^2 T^4 = L_{\odot} / (4\pi\sigma)$ (see e.g. Wittmann, 1974,). According to such a definition, the authors, by using updated model atmospheres, find that an appropriate SSM should be calculated for a radius R_{ph} smaller with respect to $R_{\odot}$ by an amount:

$$\delta R_{ph} = 0.5 \text{ Mm}; \quad (\delta R/R)_{ph} = 7 \cdot 10^{-4}. \quad (3)$$

Note that such a correction appears quite negligible in current evolutionary evaluations, achieving a relevance only in connection with the stringent constraint from HeS. As a matter of fact, in the past, SSMs (as given, e.g., by Bahcall and Pinsonneault 1995, Christensen-Dalsgaard et al. 1996, Ciacio et al. 1996) have been calculated for a radius larger than R_{ph} but, in the meantime, the suggested correction would push the radius 0.2 Mm below the value suggested by HeS.

Without entering into a discussion on that matter, in this note we discuss the effect of gravitation on the apparent solar radius. The maximum impact parameter b of photons reaching Earth corresponds to trajectories tangent to the solar surface, of radius R_0 . The relationship between b and R_0 is easily derived by using the conservation laws of energy and of angular momentum. In the weak field approximation, a photon with energy $h\nu_0$ at the Sun surface when at large distance has energy given by:

$$h\nu = h\nu_0 \left(1 - \frac{GM_{\odot}}{R_0 c^2}\right). \quad (4)$$

Angular momentum conservation gives:

$$\frac{h\nu b}{c} = \frac{h\nu_0 R_0}{c}. \quad (5)$$

The above equations immediately give:

$$b = \frac{R_0 \nu_0}{\nu} \approx R_0 + \frac{GM_{\odot}}{c^2}. \quad (6)$$

In order to get the true solar radius R_0 from the apparent solar radius $R_{\odot} = D \sin \alpha_{app} = b$ one has to subtract a quantity:

$$\delta R_{GR} = \frac{GM_{\odot}}{c^2} = 1.5 \text{ Km}; \quad (\delta R/R)_{GR} \approx 2 \cdot 10^{-6}. \quad (7)$$

This correction is small in comparison with the present accuracy of solar observations and/or helioseismic determination. However, it should be kept in mind for the future.

Acknowledgments

We appreciate discussions with J.N. Bahcall, S. Basu, E. Iacopini and W. Dziembowski. We are grateful to T. Brown, J. Christensen-Dalsgaard and G. Rybicki for pointing to our attention an important mistake in the previous version of this note.

References

- [1] Allen C.W., 1976, *Astrophysical Quantities* (3 ed.; London: Atholone Press)
- [2] Antia H.M., 1998, *A&A* 330, 336
- [3] Bahcall J.N., Pinsonneault M.H., *Rev. Mod. Phys.*, 1995, 67, 871
- [4] Bahcall J.N., Basu S., Kumar P., 1997, *ApJ* 485, L91
- [5] Basu S., 1998, in publication on *MNRAS*
- [6] Basu S., Christensen-Dalsgaard J., 1997, *A&A* 322, L5
- [7] Brown T.M., Christensen-Dalsgaard J., 1998, *ApJ* 500, L195
- [8] Ciacio F., Degl’Innocenti S., Ricci B., 1997, *A&A Suppl.* 123, 449
- [9] Christensen-Dalsgaard J. et al., 1996, *Science* 272, 1286
- [10] Degl’Innocenti S., Dziembowski W.A., Fiorentini G., Ricci B., 1997, *Astroparticle Physics* 7, 77
- [11] Dziembowski W.A., Goode P.R., Pamyatnykh A.A., Sienkiewicz R., 1994, *ApJ* 432, 417
- [12] Dziembowski W.A., 1996, *Bull. Astr. Soc. India* 24, 133
- [13] Schou J., Kosovichev A.G., Goode P.R., Dziembowski W.A., 1997, *ApJ* 489, L197
- [14] Wittmann A., 1974, *Solar Phys.* 36, 65